


How to send a referral to VPACC

Role: All providers

Step Action Activity:


1. From Add Order activity
2. Search VPACC referral by:
 - a. Type in REF1052 or use search phrase "Virtual"


Order Search

virtual 

Panels (No results found)













Medications (No results found)


Procedures 

Name	Code	Type	Priori...	F/S	Class	QI
 Referral To Virtual Pharmacist, Int - PMPA PHARMACY MED MGMT PALO ALTO	REF1052	Int Ref			PAMF Int...	

Referral To Virtual Pharmacist, Int Accept Cancel


Internal Auth • PMPA PHARMACY MED MGMT PALO ALTO, Consultation, 1 visit


Comments:       Insert SmartText      100% 


 [This referral is to the Virtual Pharmacy Ambulatory Care Clinic for optimization of treatment for heart failure patients with a clinical pharmacist.


As part of the clinical program, the pharmacist will
Review heart failure diagnosis and current and previous heart failure therapy (based on history, study


Referral:


To Department Specialty: 

To Department: 


To Provider: 


Reason: 


Type: 


Number of Visits: 

Status:

Priority: 

Scheduling Instructions:  Add Scheduling Instructions

Check-out Comments:  Add Check-out Comments

Class: 

Reference Links:

- [VPACC Collaborative Practice Agreement](#)

Accept Cancel

b. Order from Heart Failure Smart Set:

Heart Failure with Reduced EF (HFrEF) Pathway SH AMB HOD  

 [Manage User Versions](#)

This SmartSet is intended for adults with past or present reduced EF \leq 40% with prior or current HF symptoms based on the Sutter HF pathway. All 4 classes of Guideline Directed Medical Therapy (GDMT) should be used to improve outcomes HFrEF patients: (1) ARNis, ACEs, or ARBs, (2) Beta Blockers, (3) Aldosterone Antagonists, and (4) SGLT2s. Start and titrate medications to target or maximum tolerated dose, as rapidly as tolerated (within 3 months or earlier of diagnosis) to optimize outcomes.

Referral To Virtual Pharmacist, Int

Internal Auth, PMPA PHARMACY MED MGMT PALO ALTO, Consultation 1 visit